FORMATION (UPPER MIOCENE) OF CHUBUT PROVINCE, ARGENTINA, AND	
SPECIES NOT PREVIOUSLY REPORTED FROM THIS FORMATION. PART I - GAS-	
TROPODA AND SCAPHOPODA (in English and Spanish), by Rodolfo F.J. Brunet: pp.	.\$10.00
1-56 (August 30, 1995)	00.01 .
Vol. 28, Nos. 3, 4. PALEOECOLOGICAL ASPECTS OF A LATE WISCONSINAN	
AQUATIC MOLLUSK ASSEMBLAGE ASSOCIATED WITH MASTODON BONE	
RECOVERED FROM THE NORTH FORK OF WARD CREEK, BATON ROUGE,	
LOUISIANA, by Barbara Lewis; ORIGIN OF MEGAPHYTOCLAST CONCENTRA-	
TIONS IN COARSE-GRAINED TURBIDITES, CRETACEOUS OF NORTHERN	
CALIFORNIA, by William Miller, III; HOLOCENE ASTRANGIA (SCLERACTINIA) IN	
FORESHORE SHELL ACCUMULATIONS, BOGUE BANKS, NORTH CAROLINA, by	
William Miller, III; PALYNOLOGICAL CHARACTERISTICS OF NEAR-SHORE	
SHELL-BEARING PLIOCENE THROUGH HOLOCENE SEDIMENTS OF FLORIDA,	
GEORGIA, AND SOUTH CAROLINA, by Fredrick J. Rich; NOTES ON THE FAUNA	
OF THE CHIPOLA FORMATION – XXXVIII. TWO NEW SPECIES OF THE GENUS	
TURBO (GASTROPODA:TURBINIDAE), by Gary W. Schmelz; NOTES ON THE	
FAUNA OF THE CHIPOLA FORMATION – XXXIX. A NEW SPECIES OF THE	
GENUS PARAMETARIA (GASTROPODA:COLUMBELLIDAE), by Gary W. Schmelz;	
TWO ADDITIONS TO THE CENOZOIC MURICIDAE (GASTROPODA:MURICIDAE)	
OF THE WESTERN ATLANTIC REGION, by Emily H. Vokes; A NEW SPECIES OF	
ENAETA (GASTROPODA: VOLÚTIDAE) FROM THE MIO-PLIOCENE OF NORTH-	
WESTERN ECUADOR, by William D. Pitt and Lois J. Pitt; NOTES ON CERTAIN	
SPECIES OF MOLLUSCA FROM THE WACCAMAW FORMATION OF SOUTH	
CAROLINA, SAID TO BE FROM THE PINECREST FORMATION OF FLORIDA, AND	
VICE VERSA, by Richard E. Petit; ON THE OCCURRENCE OF STROMBINA	
CYPHONOTUS AND STROMBINA LESSEPSIANA IN THE DOMINICAN REPUBLIC,	
by Bernard Landau; and DISCOASTER PETALIFORMIS MOSHKOVITZ AND	
EHRLICK, 1984, COINCIDENTALIS, A NEW SUBSPECIES FROM THE SUBSUR-	
FACE OF LOUISIANA, by W.H. Akers: pp. 57-132 (November 15, 1995) - BOUND	
VOLUME 28, complete with title page and contents	.\$20.00
VOLUME 29. Subscription	.\$20.00